

Rec'd PCT/PTO 10 JUL 2004

PCT/EP 0-3. / 0.05304

21/502075



Europäisches
Patentamt

European
Patent Office

Office européen
des brevets

REC'D 24 FEB 2003

WIPO PCT

Bescheinigung

Certificate

Attestation

Die angehefteten Unterla-
gen stimmen mit der
ursprünglich eingereichten
Fassung der auf dem näch-
sten Blatt bezeichneten
europäischen Patentanmel-
dung überein.

The attached documents
are exact copies of the
European patent application
described on the following
page, as originally filed.

Les documents fixés à
cette attestation sont
conformes à la version
initialement déposée de
la demande de brevet
européen spécifiée à la
page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

02012566.2

PRIORITY DOCUMENT
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH
RULE 17.1(a) OR (b)

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

R C van Dijk



Europäisches
Patentamt

Euro
Patent Office

Office européen
des brevets

PCT/EP 03/00504

Anmeldung Nr:
Application no.: 02012566.2
Demande no:

Anmeldetag:
Date of filing: 05.06.02
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

CallistoGen AG
Neuendorfstrasse 18a
16761 Henningsdorf
ALLEMAGNE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.
If no title is shown please refer to the description.
Si aucun titre n'est indiqué se referer à la description.)

Beta-secretase inhibitors

In Anspruch genommene Priorität(en) / Priority(ies) claimed / Priorité(s)
revendiquée(s)
Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

EP/18.01.02/EP 02001339

Internationale Patentklassifikation/International Patent Classification/
Classification internationale des brevets:

C07D/

Am Anmeldetag benannte Vertragstaaten/Contracting states designated at date of
filing/Etats contractants désignées lors du dépôt:

AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU MC NL PT SE TR

05. Juni 2002

WEICKMANN & WEICKMANN

Patentanwälte

European Patent Attorneys · European Trademark Attorneys

EPO - Munich
3

05. Juni 2002

Unser Zeichen:
27196P EP-1/MDmh

Anmelder:
CallistoGen AG
Neuendorfstr. 18a

16761 Hennigsdorf

DIPL.-ING. H. WEICKMANN (bis 31.1.01)
DIPL.-ING. F. A. WEICKMANN
DIPL.-CHEM. B. HUBER
DR.-ING. H. LISKA
DIPL.-PHYS. DR. J. PRECHTEL
DIPL.-CHEM. DR. B. BÖHM
DIPL.-CHEM. DR. W. WEISS
DIPL.-PHYS. DR. J. TIESMEYER
DIPL.-PHYS. DR. M. HERZOG
DIPL.-PHYS. B. RUTTENSBERGER
DIPL.-PHYS. DR.-ING. V. JORDAN
DIPL.-CHEM. DR. M. DEY
DIPL.-FORSTW. DR. J. LACHNIT

Beta-secretase inhibitors

Beta secretase inhibitors

Description

5

The invention relates to novel beta-secretase inhibitors.

Beta-secretase (memapsin 2) is a membrane-associated aspartic protease that is involved in the production of beta-amyloid peptide. The beta-amyloid peptide is generated through proteolytic cleavage of the
10 membrane-anchored beta-amyloid precursor protein. Beta-amyloid peptide constitutes one major component of senile plaques.

Senile plaques as well as neurofibrillary tangles in the central nervous
15 tissue, however, are observed in connection with Alzheimer's disease. It is a widely accepted theory that these morphological abnormalities cause the typical cognitive malfunction (dementia) associated with Alzheimer's disease.

20 Inhibition of beta-secretase may prevent generation of senile plaques and can therefore stop the progression of Alzheimer's disease (Hong et al., Science 290 (2000) 150-153).

Therefore, much effort has been undertaken to identify beta-secretase
25 inhibitors. Inhibitors identified to date include OM991 as well as OM992, whereby the beta-secretase:OM992 complex also has been crystallized and the structure has been determined up to a resolution of 1.9Å (Ghosh et al., J.Am.Chem.Soc. 122 (2000) 3522-3523; Hong et al., Science 290 (2000) 150-153). Both, OM991 and OM992 are peptidomimetics.

30


However, since Alzheimer's disease is a wide-spread disease, with about 4 million people suffering therefrom in the U.S. alone, there is still a great need for further effective substances to treat that disease.

- 5 Therefore, it was an object of the invention to provide effective beta-secretase inhibitors.

According to the invention this object is achieved by a beta-secretase inhibitor of formula (I)

10

15




wherein

- X represents a halogen or a moiety which is bioisosteric thereto, in particular, F, Cl, Br or I, preferably Cl,
- 20 R¹ each independently represents OH or a halogen or a moiety which is bioisosteric thereto, in particular, F, Cl, Br or I, preferably Cl, and n = 0 to 4, preferably n = 0 to 2,
- R² is a connecting moiety selected from the group consisting of a single bond, a C₁-C₄ alkylene group, a C₂-C₈ alkenylene group, a C₁-C₄ alkylene group containing a least one heteroatom or C₁-C₄ alkenylene group containing at least one heteratom,
- 25 Cyc is a cyclic moiety selected from mono- and polycycles being aromatic or non-aromatic and optionally containing one or more heteroatoms selected from O, N and S, and
- 30 R³ each independently is a group being bound to the moiety Cyc and selected from halogen, NO₂, O, N, S or C₁-C₄ alkyl, C₂-C₄ alkenyl, aromatic or non-aromatic cyclic systems which, in turn, can be non-

substituted or substituted with a halogen, a OH, a C₁-C₄ alkyl or a C₁-C₄ alkenyl group, or a group selected from

5



10 and


m = 0 to 8, in particular 0 to 4.

The beta-secretase inhibitors of the invention are characterized by the presence of a halophenyl group, in particular, a chlorophenyl group, whereby a parachlorophenyl group, a diorthochlorophenyl group as well as a dimetachlorophenyl group are preferred. The phenyl group furthermore can be substituted with an OH group, with a dimetachloro-ortho-hydroxy-phenyl group being preferred.


20 The group X can be in ortho, meta or para position.

In the beta-secretase inhibitors of the invention a connecting moiety is bound to the chlorophenyl group consisting of a single bond, a C₁-C₄ alkylene group, a C₂-C₈ alkenylene group, a C₁-C₄ alkylene group containing a least one heteroatom or C₁-C₄ alkenylene group containing at least one heteroatom, preferably 1 to 3, more preferably 1 to 2 heteroatoms. Preferably, the one or more heteroatoms are selected from N, O and S, more preferably from N and S. Most preferred are connecting moieties R² containing two N atoms. The connecting moiety preferably is a single bond, a -CH₂-S-, a -CH=N-N- group or a -C(CH₃)=N-N- group.

The connecting moiety R^2 connects the halophenyl residue, in particular, a chlorophenyl residue with a further cyclic moiety. Said second cycle can be a mono- or polycycle, in particular, a polycycle condensed from two, three or four cycles. The cyclic moiety preferably contains one or more heteroatoms selected from O, N and S. Especially preferred examples of the Cyc group are




According to the invention the cyclic moiety Cyc again can be substituted with up to eight substituents, preferably up to five substituents. Examples of particularly preferred substituents on the cyclic moiety Cyc are Cl, N, methyl, allyl, paraiodophenyl, NO_2 , CF_3 as well as




Most preferably, the beta-secretase inhibitor of the invention is selected from the following compounds:


ID 1




ID 3



ID 2




ID 4





ID 5 ($IC_{50}=53,0\mu M$)

ID 6 ($IC_{50}=39,3\mu M$)



ID 7 ($IC_{50}=14,4\mu M$)



The beta-secretase inhibitors of the invention are potent compounds, by means of which beta-secretase can be inhibited selectively and effectively. They are characterized, in particular, by IC₅₀ values of $\leq 200 \mu\text{M}$. Further, the compounds of the invention provide new scaffolds for the development
5 of novel drugs based on beta-secretase inhibitors.

The compounds of the invention were identified by applying computerized screening, especially PHACIR screening, for the generation of a focussed library out of a compound data base based on a combined pharmacophore.
10 In this way it is possible to discover beta-secretase inhibitors having new structures, which had not yet been presumed in the art to have such activity.

As combined pharmacophore, for example, a common pharmacophore for
15 aspartyl proteases and a surface-based (surf2lead®) pharmacophore of the crystallized beta-secretase:OM922 complex can be used. For the common pharmacophore of the aspartyl proteases the active center was employed for generation of the pharmacophore. For the surf2lead approach the surface of the active center of the beta-secretase:OM922 complex
20 crystallized with inhibitor was used for generation of the pharmacophore. A query for PHACIR screening was generated from a combination of the two pharmacophores. The compounds of the focussed library identified by virtual screening then can be subjected to an in vitro assay, e.g. a fluorescence BACE assay, in order to thus determine possible inhibitory
25 action of the compounds.

As described above, compounds having beta-secretase inhibitory action are suitable agents for the treatment of Alzheimer's disease. The invention therefore also relates to a pharmaceutical composition comprising a beta-
30 secretase inhibitor as described above, optionally in admixture with one or more pharmaceutically acceptable carriers, diluents and/or excipients.

The pharmaceutical composition can be formulated for administration according to the respective demands. In particular, it can be formulated for topical, oral, transdermal, parenteral, inhalative or intravenous administration.

5

The amount of inventive inhibitor required for administration in the treatment and/or prophylaxis of Alzheimer's disease depends on the seriousness of the condition as well as on the patient to be treated. Typically, a daily dose is 0.01 mg/kg of body weight to 500 mg/kg of body weight, preferably at least 0.1 mg/kg of body weight to 50 mg/kg of body weight.

10

Besides the beta-secretase inhibitor the pharmaceutical compositions of the invention can contain one or more other active substances.

15

The invention further relates to the use of a beta-secretase inhibitor as described above for the manufacture of a drug for the treatment of diseases which are mediated by beta-secretase. The beta-secretase inhibitors are especially suited for the production of a drug for the treatment of Alzheimer's disease. The expression "treatment of a condition" as used herein refers both to the treatment of established symptoms and a prophylactic treatment, by which the occurrence of the disease or particular symptoms can be avoided.

20

25 The invention is further illustrated by the following Example.

Example 1

Fluorescence BACE assay

30

The inhibitory activity of the compounds of the invention was shown in an in vitro assay, namely a fluorescence BACE assay.

The assay was set up in triplicate wells of 96 well black plate. rhBACE was diluted to 1 unit/well in 100 μ l (PBS + 0.5% Triton-X 100, pH5). BACE enzyme (obtained from R&D systems (ca.No.931-AS), reference: Vasser et al., 1999, Science 286, 735-741) was incubated with various concentrations of inhibitor compound (10 nM to 500 μ M) for 5 min. Reaction was started by adding peptide substrate (obtained from BACHEM (cat. No.M-2470), reference: Ermolieff et al., Biochemistry 39 (2000) 12450-56) with EDANS/Dabcyl labels. After incubation for 2 hours at 37°C the results were read in fluoroplate reader at 355 nm/486 nm.

The following IC50 values were determined for the above-mentioned particularly preferred compounds:

ID1: IC50 = 38 μ M; ID2: IC50 = 52 μ M; ID3: IC50 = 90 μ M; ID4: IC50 = 140-170 μ M; ID5: IC50 = 53 μ M; ID6: IC50 = 39.3 μ M and ID7: IC50 = 14.4 μ M.

Claims


EPO - Munich
3

05. Juni 2002

1. Beta-secretase inhibitor of formula (I)

5


10



wherein

- X represents a halogen or a moiety which is bioisosteric thereto,
- R¹ each independently represents OH or a halogen or a moiety which is bioisosteric thereto, and n = 0 to 4,
- 15 R² is a connecting moiety selected from the group consisting of a single bond, a C₁-C₄ alkylene group, a C₂-C₈ alkenylene group, a C₁-C₄ alkylene group containing a least one heteroatom or a C₁-C₄ alkenylene group containing at least one heteratom,
- 20 Cyc is a cyclic moiety selected from monocycles and polycycles being aromatic or non-aromatic and optionally containing one or more heteroatoms selected from O, N and S, and
- R³ each independently is a group being bound to the moiety Cyc and selected from halogen, O, N, S, NO₂ or C₁-C₄ alkyl, C₂-C₄ alkenyl, O, N, S, aromatic or non-aromatic cyclic systems which, in turn, can be non-substituted or substituted with halogen, OH, a C₁-C₄ alkyl or a C₁-C₄ alkenyl group, or a group selected from

30




and
m = 0 to 8.

2. Beta-secretase inhibitor according to claim 1 having the formula


5

ID 1




10

ID 3




15

ID 2




20

ID 4




ID 5 ($IC_{50}=53,0\mu M$)



25


30

ID 6 ($IC_{50}=39,3\mu M$)



ID 7 ($IC_{50}=14,4\mu M$)

or



3. Beta-secretase inhibitor according to claim 1 or 2,
having an $IC_{50} \leq 200 \mu M$.
4. A pharmaceutical composition comprising a beta-secretase inhibitor
according to any of claims 1 to 3,
optionally in admixture with one or more pharmaceutically
acceptable carriers, diluents and/or excipients.
5. The use of a beta-secretase inhibitor according to any of claims 1 to
3 for the manufacture of a pharmaceutical agent for the treatment of
a condition which is mediated by beta-secretase.
6. The use according to claim 5 for the manufacture of a
pharmaceutical agent for the treatment of Alzheimer's disease.

05. Juni 20

- 12 -

Abstract

EPO - Munich
3

05. Juni 2002

The invention relates to novel beta-secretase inhibitors.

5

mh 04.06.2002